Ensemble Ramachandran plots
SR213_NMR_em_bcr3 (20 models)**



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

** Selected residues only

Ensemble Ramachandran plots
SR213_NMR_em_bcr3 (20 models)**



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

** Selected residues only

Ensemble Ramachandran plots
SR213_NMR_em_bcr3 (20 models)**



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

** Selected residues only

Ensemble Ramachandran plots
SR213_NMR_em_bcr3 (20 models)**



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

** Selected residues only

Ensemble Ramachandran plots
SR213_NMR_em_bcr3 (20 models)**



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

** Selected residues only